

Predictions for deeply virtual Compton scattering on a spin-one target

Axel Kirchner¹ and Dieter Müller²

¹ Institut für Theoretische Physik, Universität Regensburg, D-93040 Regensburg, Germany

² Fachbereich Physik, Universität Wuppertal, D-42097 Wuppertal, Germany

Abstract. We consider hard leptonproduction of a photon on a spin-one target and give the general azimuthal angular dependence of the differential cross section. Furthermore, we estimate the beam spin asymmetry for an unpolarized deuteron target at HERMES.

1 Introduction

Exclusive two-photon processes in the light-cone dominated region, i.e., in the generalized Bjorken limit, are most suitable for the exploration of the partonic content in hadrons, since in leading order (LO) both photons directly couple to one quark line [1]. In such processes one can measure, for instance, different photon-to-meson form factors, i.e., $\gamma^*\gamma \rightarrow M$, the production of hadron pairs and also processes like $\gamma^*N \rightarrow N\gamma$ or $\gamma N \rightarrow Nl^+l^-$. The latter two are denoted as deeply virtual Compton scattering (DVCS) in the space- and time-like regions [2,3,4,5,6,7].

The factorization of short- and long-range dynamics is formally given by the operator product expansion (OPE) of the time ordered product of two electromagnetic currents, which has been worked out at leading twist-two in next-to-leading order (NLO) and at twist-three level in LO of perturbation theory (for references see [7]). However, one should be aware that the partonic hard-scattering part, i.e., the Wilson coefficients, contains collinear singularities, which are absorbed in the non-perturbative distributions by a factorization procedure, which has been proven at twist-two level [8].

The non-perturbative distributions are defined in terms of light-ray operators with definite twist sandwiched between the corresponding hadronic states. These process dependent correlation functions are sensitive to different aspects of hadronic physics. Especially, in DVCS one can access the so-called generalized parton distributions (GPDs). The second moment of the flavour singlet GPDs is related to the expectation value of the energy momentum tensor. Thus, it gives in principle information on the angular orbital momentum fraction of the nucleon spin carried by quarks [9]. We should stress that this process is a new tool to probe the partonic content of the nucleon on the level of amplitudes and, thus, it provides us new information [10].

Recently, the DVCS process has been measured by the H1 collaboration [11] in the small x_B region (see also [12]) as well as in single beam spin asymmetries by the HERMES [13] and CLAS [14] collaborations. The theoretical predictions

depend on the GPDs, which are unknown. However, they are constrained by sum rules and the reduction to the parton densities in the forward kinematics. All experimental data are consistent with an oversimplified model, which satisfies the constraints and, thus, also with each other [7].

It is appealing to employ DVCS for the investigation of other hadrons and nuclei. An appropriate candidate is deuteron, which has been widely used as a target in lepto-scattering experiments. This nucleus has been extensively studied in both deep-inelastic [15] and elastic [16,17] scattering. From the theoretical point of view it would be desired to have complementary information, which could give us a deeper understanding of this nucleus in terms of its fundamental degree of freedom [18].

In this paper we outline the OPE approach applied to DVCS. Moreover, we determine the azimuthal angular dependence of the cross section for a spin-1 target. Relying on qualitative properties of GPDs, which are consistent with the DVCS data for the proton target, we estimate the size of the beam spin asymmetry for HERMES kinematics.

2 General formalism

For the leptonproduction of a photon

$$l^\pm(k)h(P_1) \rightarrow l^\pm(k')h(P_2)\gamma(q_2) \quad (1)$$

on a hadronic target h with the mass M the five-fold cross section

$$\frac{d\sigma}{dx_B dy d|\Delta^2| d\phi d\varphi} = \frac{\alpha^3 x_B y}{16 \pi^2 Q^2 \sqrt{1 + \epsilon^2}} \left| \frac{\mathcal{T}}{e^3} \right|^2, \quad \epsilon \equiv 2x_B \frac{M}{Q}, \quad (2)$$

depends on the Bjorken variable $x_B = Q^2/2P_1 \cdot q_1$, where $Q^2 = -q_1^2$ with $q_1 = k - k'$, the momentum transfer square $\Delta^2 = (P_2 - P_1)^2$, the lepton energy fraction $y = P_1 \cdot q_1 / P_1 \cdot k$ and, in general, two azimuthal angle. We use the target rest frame, where the virtual photon three-momentum points towards the negative z -direction. ϕ is the angle between the lepton and hadron scattering planes and $\varphi = \Phi - \phi_N$ is the difference of the azimuthal angle Φ of the spin vector

$$S^\mu = (0, \cos \Phi \sin \Theta, \sin \Phi \sin \Theta, \cos \Theta) \quad (3)$$

and the azimuthal angle ϕ_N of the recoiled hadron (see Ref. [7]).

In the following we consider this process in the (generalized) Bjorken limit, $Q^2 \sim P_1 \cdot q_1$ is large, Δ^2 and M^2 are comparably small. The amplitude $\mathcal{T}$ is the sum of the DVCS $\mathcal{T}_{\text{DVCS}}$ and Bethe-Heitler (BH) $\mathcal{T}_{\text{BH}}$ amplitudes:

$$\mathcal{T}^2 = \sum_{\lambda', \lambda'} \{ |\mathcal{T}_{\text{BH}}|^2 + |\mathcal{T}_{\text{DVCS}}|^2 + \mathcal{I} \}, \quad (4)$$

with the interference term

$$\mathcal{I} = \mathcal{T}_{\text{DVCS}} \mathcal{T}_{\text{BH}}^* + \mathcal{T}_{\text{DVCS}}^* \mathcal{T}_{\text{BH}}, \quad (5)$$

where the recoiled lepton (λ') and hadron (A') polarization will not be observed.

Each of these three terms in Eq. (4) can be calculated in a straightforward manner by the contraction of a known leptonic tensor $L_{\mu\dots}$ with the DVCS tensor $T_{\mu\nu}^{\text{DVCS}}$ or/and hadronic current J_α . The former one is given in terms of GPDs, while the latter one is parametrized in terms of elastic electromagnetic form factors. For instance, the interference term reads

$$\sum_{\lambda'} \mathcal{I} = \frac{\pm e^6}{Q^2 \Delta^2} L^{\mu\nu\alpha}(k, k') T_{\mu\nu}^{\text{DVCS}}(P, \Delta, q) J_\alpha^\dagger(\Delta) + \text{h.c.} \quad \begin{cases} + & \text{for } e^- \\ - & \text{for } e^+ \end{cases}, \quad (6)$$

where $P = P_1 + P_2$ and $q = (q_1 + q_2)/2$. The resulting predictions for the spin-0 and -1/2 targets are presented at the twist-three level in Refs. [5,7].

Let us now consider the spin-1 target in more detail. The hadronic current

$$J_\mu = -\epsilon_2^* \cdot \epsilon_1 P_\mu G_1 + (\epsilon_2^* \cdot P \epsilon_{1\mu} + \epsilon_1 \cdot P \epsilon_{2\mu}^*) G_2 - \epsilon_2^* \cdot P \epsilon_1 \cdot P \frac{P_\mu}{2M^2} G_3 \quad (7)$$

is given by three form factors $G_i(\Delta^2)$ with $i = \{1, 2, 3\}$, where $\epsilon_{1\mu}$ ($\epsilon_{2\mu}$) denote the three polarization vectors for the initial (final) hadron. The form factors $G_i(\Delta^2)$ have been measured and their parametrizations are available in the literature, see Ref. [17] and references therein.

The DVCS hadronic tensor is given by the time-ordered product of the electromagnetic currents j_μ , which is sandwiched between hadronic states with different momenta. In LO of perturbation theory and at twist-two accuracy it reads [4]

$$\begin{aligned} T_{\mu\nu}(\xi, \Delta^2, Q^2) &= \frac{i}{e^2} \int dx e^{ix \cdot q} \langle P_2 | T j_\mu(x/2) j_\nu(-x/2) | P_1 \rangle \\ &= -\mathcal{P}_{\mu\sigma} g_{\sigma\tau} \mathcal{P}_{\tau\nu} \frac{q \cdot V_1}{P \cdot q} - \mathcal{P}_{\mu\sigma} i \epsilon_{\sigma\tau q\rho} \mathcal{P}_{\tau\nu} \frac{A_1 \rho}{P \cdot q}, \end{aligned} \quad (8)$$

where the projection operator $\mathcal{P}_{\mu\nu} = g_{\mu\nu} - q_{1\mu} q_{2\nu} / q_1 \cdot q_2$ ensures gauge invariance. For convenience we introduced the scaling variable $\xi \approx \frac{x_B}{2-x_B}$. At twist-two level the amplitudes V_1 and A_1 can be decomposed in a complete basis of nine Compton form factors (CFFs). Adopting the notation of Ref. [18], they read in the vector case

$$\begin{aligned} V_\mu &= -\epsilon_2^* \cdot \epsilon_1 P_\mu \mathcal{H}_1 + (\epsilon_2^* \cdot P \epsilon_{1\mu} + \epsilon_1 \cdot P \epsilon_{2\mu}^*) \mathcal{H}_2 - \epsilon_2^* \cdot P \epsilon_1 \cdot P \frac{P_\mu}{2M^2} \mathcal{H}_3 \\ &\quad + (\epsilon_2^* \cdot P \epsilon_{1\mu} - \epsilon_1 \cdot P \epsilon_{2\mu}^*) \mathcal{H}_4 + \left(\frac{2M^2 \{ \epsilon_2^* \cdot q \epsilon_{1\mu} + \epsilon_1 \cdot q \epsilon_{2\mu}^* \}}{P \cdot q} + \frac{\epsilon_2^* \cdot \epsilon_1}{3} P_\mu \right) \mathcal{H}_5, \end{aligned} \quad (9)$$

and in the axial-vector case

$$\begin{aligned} A_\mu &= i \epsilon_{\mu\epsilon_2^* \epsilon_1 P} \tilde{\mathcal{H}}_1 - \frac{i \epsilon_{\mu\Delta P \epsilon_1} \epsilon_2^* \cdot P + i \epsilon_{\mu\Delta P \epsilon_2^*} \epsilon_1 \cdot P}{M^2} \tilde{\mathcal{H}}_2 \\ &\quad - \frac{i \epsilon_{\mu\Delta P \epsilon_1} \epsilon_2^* \cdot P - i \epsilon_{\mu\Delta P \epsilon_2^*} \epsilon_1 \cdot P}{M^2} \tilde{\mathcal{H}}_3 - \frac{i \epsilon_{\mu\Delta P \epsilon_1} \epsilon_2^* \cdot q + i \epsilon_{\mu\Delta P \epsilon_2^*} \epsilon_1 \cdot q}{q \cdot P} \tilde{\mathcal{H}}_4, \end{aligned} \quad (10)$$

where $1/Q$ -power suppressed effects have been neglected. The remaining logarithmic Q -dependence is governed by perturbation theory.

The CFFs in Eqs. (9) and (10) are given by a convolution of perturbatively calculable coefficient functions $C^{(\pm)}$ and twist-two GPDs via

$$\mathcal{H}_k(\xi) = \sum_{i=u,\dots} \int_{-1}^1 dx C_i^{(-)}(\xi, x) H_k^i(x, \eta)|_{\eta=-\xi}, \quad \text{for } k = \{1, \dots, 5\}, \quad (11)$$

$$\tilde{\mathcal{H}}_k(\xi) = \sum_{i=u,\dots} \int_{-1}^1 dx C_i^{(+)}(\xi, x) \tilde{H}_k^i(x, \eta)|_{\eta=-\xi}, \quad \text{for } k = \{1, \dots, 4\}. \quad (12)$$

For each quark species i we have nine GPDs. The two sets $\{H_1^i, \dots, H_5^i\}$ and $\{\tilde{H}_1^i, \dots, \tilde{H}_4^i\}$ are defined by off-forward matrix elements of vector and axial-vector light-ray operators. The coefficient functions $C^{(\mp)}$ have perturbative expansion. In LO they read for the even $(-)$ and odd $(+)$ parity sectors

$$\xi C_{(0)i}^{(\mp)}(\xi, x) = \frac{Q_i^2}{1 - x/\xi - i0} \mp \frac{Q_i^2}{1 + x/\xi - i0}, \quad (13)$$

where Q_i is the fractional quark charge.

Employing the parametrizations (7), (9) and (10), the contractions of leptonic and hadronic tensors provide the kinematically exact expression for the squared BH term (of course, in tree approximation), while the interference term

$$\begin{aligned} \sum_{\lambda'} \mathcal{T}_{\text{BH}} \mathcal{T}_{\text{DVCS}}^* = & \quad (14) \\ & \frac{2 - 2y + y^2}{y^2 \mathcal{P}_1(\phi) \mathcal{P}_2(\phi)} \frac{4\xi}{\Delta^2 Q^4} \left(k^\sigma - \frac{q^\sigma}{y} \right) \left[\left(J_\sigma + 2\Delta_\sigma \frac{q \cdot J}{Q^2} \right) q \cdot V_1^\dagger + 2i\epsilon_{\sigma q \Delta J} \frac{q \cdot A_1^\dagger}{Q^2} \right] \\ & + \frac{\lambda(2-y)y}{y^2 \mathcal{P}_1(\phi) \mathcal{P}_2(\phi)} \frac{4\xi}{\Delta^2 Q^4} \left(k^\sigma - \frac{q^\sigma}{y} \right) \left[\left(J_\sigma + 2\Delta_\sigma \frac{q \cdot J}{Q^2} \right) q \cdot A_1^\dagger + 2i\epsilon_{\sigma q \Delta J} \frac{q \cdot V_1^\dagger}{Q^2} \right], \end{aligned}$$

and the squared DVCS amplitude

$$\begin{aligned} \sum_{\lambda'} |\mathcal{T}_{\text{DVCS}}|^2 = & 8 \frac{2 - 2y + y^2}{y^2} \frac{\xi^2}{Q^6} \left(q \cdot V_1 \ q \cdot V_1^\dagger + q \cdot A_1 \ q \cdot A_1^\dagger \right) \\ & + 8 \frac{\lambda(2-y)}{y} \frac{\xi^2}{Q^6} \left(q \cdot V_1 \ q \cdot A_1^\dagger + q \cdot A_1 \ q \cdot V_1^\dagger \right) \end{aligned} \quad (15)$$

are expanded with respect to $1/Q$. In contrast to the squared DVCS amplitude the interference as well as the squared BH terms have an additional ϕ -dependence due to the (scaled) BH propagators

$$\mathcal{P}_1 \approx -\frac{1}{y} \{1 - y + 2K \cos(\phi)\}, \quad \mathcal{P}_2 \approx \frac{1}{y} \{1 + 2K \cos(\phi)\}. \quad (16)$$

The kinematical factor

$$K \approx \sqrt{-\frac{\Delta^2}{Q^2}(1-x_B)(1-y)} \left(1 - \frac{\Delta_{\min}^2}{\Delta^2}\right). \quad (17)$$

vanishes at the kinematical boundary $\Delta^2 = \Delta_{\min}^2 \approx M^2 x_B^2 / (1-x_B)$.

3 Observables for leptonproduction of a photon

In our frame the contractions of leptonic and hadronic tensors, see Eq. (6), yield finite sums of Fourier harmonics, whose form is governed by general principles. After summation over the final polarization states, which is not indicated below, the three parts of the squared amplitude read for massless leptons [7]:

$$|\mathcal{T}_{\text{BH}}|^2 = \frac{e^6(1+\epsilon^2)^{-2}}{x_B^2 y^2 \Delta^2 \mathcal{P}_1(\phi) \mathcal{P}_2(\phi)} \left\{ c_0^{\text{BH}} + \sum_{n=1}^2 [c_n^{\text{BH}} \cos(n\phi) + s_n^{\text{BH}} \sin(\phi)] \right\}, \quad (18)$$

$$|\mathcal{T}_{\text{DVCS}}|^2 = \frac{e^6}{y^2 Q^2} \left\{ c_0^{\text{DVCS}} + \sum_{n=1}^2 [c_n^{\text{DVCS}} \cos(n\phi) + s_n^{\text{DVCS}} \sin(n\phi)] \right\}, \quad (19)$$

$$\mathcal{I} = \frac{\pm e^6}{x_B y^3 \mathcal{P}_1(\phi) \mathcal{P}_2(\phi) \Delta^2} \left\{ c_0^{\mathcal{I}} + \sum_{n=1}^3 [c_n^{\mathcal{I}} \cos(n\phi) + s_n^{\mathcal{I}} \sin(n\phi)] \right\}, \quad (20)$$

where the $+$ ($-$) sign in the interference stands for the negatively (positively) charged lepton beam.

The Fourier coefficients can be calculated from Eqs. (14), (15), and an analogous one for the squared BH amplitude by summing over the polarization Λ' , where we can employ the common projector technique. For a spin-1 particle we have (see for instance [19])

$$\begin{aligned} \epsilon_{1\mu}^*(\Lambda=0)\epsilon_{1\nu}(\Lambda=0) &= S_\mu S_\nu, \\ \epsilon_{1\mu}^*(\Lambda=\pm 1)\epsilon_{1\nu}(\Lambda=\pm 1) &= \frac{1}{2} \left(-g_{\mu\nu} + \frac{P_{1\mu} P_{1\nu}}{M^2} - S_\mu S_\nu + \frac{i\Lambda}{M} \epsilon_{\mu\nu P_1 S} \right), \end{aligned} \quad (21)$$

where $\Lambda = \{+1, 0, -1\}$ denotes the magnetic quantum number with respect to the quantization direction given by the spin vector S_μ defined in Eq. (3). Obviously, the spin sum of the recoiled hadron is

$$\sum_{\Lambda'=-1}^1 \epsilon_{2\mu}^*(\Lambda')\epsilon_{2\nu}(\Lambda') = -g_{\mu\nu} + \frac{P_{2\mu} P_{2\nu}}{M^2}. \quad (22)$$

As we see, for a spin-1 target the Fourier coefficients quadratically depend on the spin vector S_μ and, thus, we find the following decomposition

$$\begin{aligned} c_n^T &= c_{n,\text{unp}}^T + c_{n,\text{LP}}^T \cos(\Theta) + c_{n,\text{TP}}^T(\varphi) \sin(\Theta) + c_{n,\text{LTP}}^T(\varphi) \sin(2\Theta) \\ &\quad + c_{n,\text{LLP}}^T \cos^2(\Theta) + c_{n,\text{TTP}}^T(\varphi) \sin^2(\Theta) \end{aligned} \quad (23)$$

for $T = \{\text{BH}, \mathcal{I}, \text{DVCS}\}$. An analogous decomposition holds true for the odd harmonics s_n^T . The unpolarized and the longitudinally polarized coefficients $c/s_{n,\text{unp}}^T$, $c/s_{n,\text{LP}}^T$, and $c/s_{n,\text{LLP}}^T$, respectively, are independent of φ . The transverse coefficients $c/s_{n,\text{TP}}^T$ and the transverse-longitudinal interference terms can be decomposed with respect to the first harmonics in the azimuthal angle φ

$$c_{n,\text{TP}}^T(\varphi) = c_{n,\text{TP}+}^T \cos(\varphi) + s_{n,\text{TP}-}^T \sin(\varphi), \quad (24)$$

$$c_{n,\text{LTP}}^T(\varphi) = c_{n,\text{LTP}+}^T \cos(\varphi) + s_{n,\text{LTP}-}^T \sin(\varphi), \quad (25)$$

while $c/s_{n,\text{TTP}}^T$ may be written as

$$c_{n,\text{TTP}}^T(\varphi) = c_{n,\text{TTP}\Sigma}^T + c_{n,\text{TTP}\Delta}^T \cos(2\varphi) + s_{n,\text{TTP}\pm}^T \sin(2\varphi). \quad (26)$$

Analogous equations hold true for the odd harmonics (just replace $c \leftrightarrow s$). Let us add that with this notation the $c(s)$ harmonics are given by the real (imaginary) part of certain linear combinations of form factors and/or CFFs.

Altogether we have for a given harmonic in ϕ nine possible observables¹. In principle, they can be measured by an appropriate adjustment of the spin vector and Fourier analysis with respect to the azimuthal angle φ . The interference term linearly depends on the CFFs and is, thus, of special interest. In facilities that have both kinds of leptons it can be separated by means of the charge asymmetry. The dominant harmonics are c/s_1^T , predicted at leading twist-two. We write them as product of leptonic prefactors $\mathcal{L}$ and ‘universal’ functions $\mathcal{C}_i^T$:

$$\begin{Bmatrix} c_{1,i}^{\mathcal{I}} \\ s_{1,i}^{\mathcal{I}} \end{Bmatrix} = \begin{Bmatrix} \mathcal{L}_{1,i}^{\mathcal{I}c} \\ \mathcal{L}_{1,i}^{\mathcal{I}s} \end{Bmatrix} \begin{Bmatrix} \Re \\ \Im \end{Bmatrix} \mathcal{C}_i^T, \quad \text{for } i = \{\text{unp}, \dots, \text{TTP-}\}. \quad (27)$$

As mentioned before, they depend on nine CFFs

$$\mathcal{C}_i^T = (G_1 \cdots G_3) \mathcal{M}_i^T \begin{pmatrix} \mathcal{H}_1 \\ \vdots \\ \tilde{\mathcal{H}}_4 \end{pmatrix}, \quad (28)$$

where the matrix $\mathcal{M}$ will be presented elsewhere [20]. Single spin-flip and unpolarized as well as double spin-flip contributions provide the imaginary and real part, respectively, of the nine linear combinations $\mathcal{C}_i^T$.

4 Estimate of the beam-spin asymmetry

In this section we estimate the size of the beam spin asymmetry

$$A_{\text{LU}}(\phi) = \frac{d\sigma^\uparrow(\phi) - d\sigma^\downarrow(\phi)}{d\sigma^\uparrow(\phi) + d\sigma^\downarrow(\phi)}, \quad (29)$$

¹ Note that $c_{n,\text{TTP}\Sigma}^T$ does not belong to an independent frequency, rather it can be included in the constant and $\cos^2(\theta)$ terms of Eq. (23).

at large y and small momentum transfer for the HERMES experiment. We should also ensure that M^2/Q^2 corrections are under control. Since $M = 1875.6$ MeV we require $Q^2 \geq 4$ GeV². Because of large y , the BH amplitude dominates the DVCS one and thus the beam spin asymmetry is approximately determined by

$$A_{\text{LU}}(\phi) \sim \pm \frac{x_{\text{B}} s_{1,\text{unp}}^{\mathcal{I}}}{y c_{0,\text{unp}}^{\text{BH}}} \sin(\phi) \quad \text{with} \quad \begin{cases} + & \text{for } e^- \\ - & \text{for } e^+ \end{cases}. \quad (30)$$

For $-\Delta_{\text{min}} \ll -\Delta^2 \ll M^2$ and $x_{\text{B}} \lesssim 0.3$ the Fourier coefficients can be drastically simplified due a crude approximation of kinematical factors

$$A_{\text{LU}}(\phi) \sim \pm \sqrt{\frac{-\Delta^2}{Q^2}(1-y)} \frac{x_{\text{B}} \Im \mathcal{H}_1(\xi, \Delta^2, Q^2)|_{\xi=x_{\text{B}}/2}}{G_1(\Delta^2)} \sin(\phi). \quad (31)$$

For a spin-1/2 target one finds the analogous equation, i.e., $G_1 \rightarrow F_1$ and $\mathcal{H}_1 \rightarrow \mathcal{H}$. Note that the same rule

$$\int_{-1}^1 dx \left(Q_u \left\{ \frac{H_1}{H} \right\}_{u_v}(x, \xi, \Delta^2, Q^2) + Q_d \left\{ \frac{H_1}{H} \right\}_{d_v}(x, \xi, \Delta^2, Q^2) \right) = \left\{ \frac{G_1}{F_1} \right\}(\Delta^2) \quad (32)$$

suggest that the Δ^2 -dependence of the valence-quark GPDs is essentially given by G_1 and F_1 , respectively. The analyses of the H1, HERMES and CLAS data at LO indicate that the Δ^2 fall-off of the sea-quark GPDs is steeper than the valence quark ones. Thus, we neglect the sea quark contribution and write the valence like GPDs as a product of form factor and quark distribution function

$$\frac{\Im \mathcal{H}(\xi, \Delta^2)}{F_1(\Delta^2)} \sim \pi \{ Q_u^2 q_{u_v}(\xi) + Q_d^2 q_{d_v}(\xi) \}_{|\xi=x_{\text{B}}/2}, \quad (33)$$

$$\frac{\Im \mathcal{H}_1(\xi, \Delta^2)}{G_1(\Delta^2)} \sim \pi \frac{Q_u^2 + Q_d^2}{2} \{ q_{u_v}(\xi) + q_{d_v}(\xi) \}_{|\xi=x_{\text{B}}/2}. \quad (34)$$

For HERMES kinematics with central values $\langle x_{\text{B}} \rangle = 0.11$, $\langle \Delta^2 \rangle = -0.27$ GeV² and $\langle Q^2 \rangle = 2.6$ GeV² we find with the MRS A' parametrization $A_{\text{LU}}(\phi) \sim 0.3 \sin(\phi)$ for positron-proton scattering, which is consistent with experimental data [13]. For $Q^2 = 4$ GeV², $x_{\text{B}} = 0.1$ and $\Delta^2 = -0.3$ GeV² we estimate

$$\begin{aligned} A_{\text{LU}}(\phi) &\sim -0.13 \sin(\phi) && \text{for deuteron target,} \\ A_{\text{LU}}(\phi) &\sim -0.16 \sin(\phi) && \text{for proton target.} \end{aligned} \quad (35)$$

5 Summary

We gave the general azimuthal angular dependence of the leptonproduction cross section of a photon on a spin-1 target. Such experiments allow to study the deuteron from a new perspective. Compared to DIS or elastic lepton-deuteron scattering, they provide additional information, contained in the CFFs.

At twist-two level there are nine CFFs. In the case of a polarized beam and target, with an adjustable quantization direction, the imaginary and real part of all these CFFs can be separately measured by means of the charge asymmetry. Moreover, an appropriate Fourier analysis allows to eliminate the twist-three contamination. In this way one gets the maximal access to the deuteron GPDs at twist-two level, with a contamination of M^2/Q^2 and Δ^2/Q^2 contributions.

As discussed, for certain kinematics the beam spin asymmetry is essentially determined by one CFF only. Although the approximations are rather crude, the result can serve to obtain a qualitative understanding of this GPD. The LO analysis of the pioneering measurements of DVCS [11,13,14] on the proton suggest for $-\Delta^2 \sim 0.3 \text{ GeV}^2$ the dominance of valence-quark GPDs with no essential enhancement by the skewedness effect. Assuming the same features also for the deuteron GPDs, we estimated the size of the beam spin asymmetry.

References

1. D. Müller, D. Robaschik, B. Geyer, F.-M. Dittes, and J. Hořejší, Fortschr. Phys. **42**, 101 (1994).
2. A. Radyushkin, Phys. Lett. B **380**, 417 (1996); X. Ji, Phys. Rev. D **55**, 7114 (1997).
3. M. Diehl, T. Gousset, B. Pire, and J. P. Ralston, Phys. Lett. B **411**, 193 (1997).
4. A. Belitsky, D. Müller, L. Niedermeier, and A. Schäfer, Nucl. Phys. B **593**, 289 (2001).
5. A. Belitsky, D. Müller, A. Kirchner, and A. Schäfer, Phys. Rev. D **64**, 116002 (2001).
6. E. R. Berger, M. Diehl, and B. Pire, *Timelike Compton scattering: Exclusive photoproduction of lepton pairs*, hep-ph/0110062 (2001).
7. A. V. Belitsky, D. Müller, and A. Kirchner, *Theory of deeply virtual compton scattering on the nucleon*, hep-ph/0112108 (2001).
8. J. Collins and A. Freund, Phys. Rev. D **59**, 074009 (1999); X. Ji and J. Osborne, Phys. Rev. D **58**, 094018 (1998).
9. X. Ji, Phys. Rev. Lett. **78**, 610 (1997).
10. J. Ralston and B. Pire, *Femto-photography of protons to nuclei with deeply virtual Compton scattering*, (2001).
11. C. Adloff et al., Phys. Lett. B **517**, 47 (2001).
12. P. R.B. Saull (ZEUS Coll.), *Prompt photon production and observation of deeply virtual compton scattering*, hep-ex/0003030 (2000).
13. A. Airapetian et al., Phys. Rev. Lett. **87**, 182001 (2001).
14. S. Stepanyan et al., Phys. Rev. Lett. **87**, 182002 (2001).
15. P. Hoodbhoy, R. Jaffe, and A. Manohar, Nucl. Phys. B **312**, 571 (1989).
16. S. Brodsky, C.-R. Ji, and G. Lepage, Phys. Rev. Lett. **51**, 83 (1983).
17. R. Gilman and F. Gross, *Electromagnetic structure of the deuteron*, nuc-th/0111015 (2001); M. Garçon and J. V. Orden, *The deuteron: structure and form factors*, nuc-th/0102049 (2001).
18. E. R. Berger, F. Cano, M. Diehl, and B. Pire, Phys. Rev. Lett. **87**, 142302 (2001).
19. G. Passarino, Nucl. Phys. B **237**, 249 (1984).
20. A. Kirchner and D. Müller, *in preparation*.